


Meyer, A. $^1$ , Happold, J. $^1$ , Sadler R. $^1$ , Zalcman, E. $^1$ , Hillman, A. $^1$ , Mackenzie, C. $^1$ , Cowled, B. $^1$ , Gallardo Lagno, A.L. $^2$ , Stegeman, J.A. $^3$ , Cameron, A. $^1$ 

Salmonid rickettsial septicaemia
Caused by the bacterium
Piscirickettsia salmonis, SRS is the
major cause for antibiotic
treatment of salmon and trout in
Chile.



## Methods used in this PhD project:

- Stakeholder consultations
- Creation of a platform integrating health and management data from individual companies
- Epidemiological studies



Lessons learned: building relationships with stakeholders takes time and effort but it is crucial for the success of data

- Data harmonisation is a complex and time-consuming task requiring stakeholder inputs on a regular basis
- Stakeholder consultations contribute to better understand a complex system before conducting in-depth data analysis
- Providing outputs that can be used to support decisionmaking is critical to maintain engagement, improve knowledge transfer and increase the impact of the work





University of Western Australia, M089 School of Agriculture and Environment, 35 Stirling Hwy, Crawley, WA 6008, Australia



What are the effects of lice burden

and lice bathing treatments on SRS







integration:

<sup>1</sup> Ausvet Europe, Lyon, France and Ausvet, Bruce, Australia <sup>2</sup> Servicio Nacional de Pesca y Acuicultura, Valparaíso, Chile <sup>3</sup> Department of Farm Animal Health of Utrecht University, Utrecht, the Netherlands